

## Adolescent Nutrition 3



## Strategies and interventions for healthy adolescent growth, nutrition, and development

Dougal Hargreaves\*, Emily Mates\*, Purnima Menon\*, Harold Alderman, Delan Devakumar, Wafai Fawzi, Geva Greenfield, Weeam Hammoudeh, Shanshan He, Anwasha Lahiri, Zheng Liu, Phuong Hong Nguyen, Vani Sethi, Haijun Wang, Lynnette M Neufeld†, George C Patton†

Adolescence is a pivotal point in the life course, characterised by transformative physical, cognitive, and emotional growth, an openness to change, and a drive to reshape the social environment. It offers unique opportunities to adopt changes in diet and physical activity that can persist into later life. Yet pre-existing nutritional problems, including micronutrient deficiencies, food insecurity, and poor-quality diets, persist at the same time as adolescents face the rapid emergence of an obesity epidemic. Adolescent growth and nutrition has been largely overlooked in intervention and policy research. Most intervention studies have emphasised micronutrient supplementation, with few taking into account the multiple drivers of adolescent diets. This Series paper highlights that effective interventions and policies will need to cut across sectors; be supported by multifaceted and multilevel policy; and extend across education, health, food systems, social protection, and digital media. Better data standardisation and systems will be essential in coordinating and monitoring these responses. In a context of shifts in planetary ecosystems and commercial drivers, resilient food systems will need to both ensure access to healthy and affordable foods and provide the infrastructure and incentives for continuing physical activity. Intergenerational partnerships with young people will be essential in bringing about transformative change and ensuring that food policies reflect their needs and aspirations.

## Introduction

Adolescence (10–24 years of age) is characterised by transition, exploration, and openness to change, offering opportunities for radical shifts in diet, physical activity, and other risks for non-communicable diseases. This same novelty-seeking and openness to change also makes adolescents a vulnerable group to commercial exploitation and other unhealthy influences, with lifelong and intergenerational consequences. Given the rapid and transformative growth of puberty, adolescents' nutritional requirements differ from both younger and older groups.<sup>1</sup> Yet, even as nutrition has become more prominent in the nurturing of human capital, adolescents have still not been included in the global nutritional policies and frameworks.<sup>2</sup> The consequences have been a failure to invest consistently in either adolescent-responsive nutritional programmes or research. There has been too much focus on single micronutrient supplements and a failure to tackle the multiple drivers of malnutrition and poor diet.<sup>3</sup> As a result, long-standing nutritional problems such as iron deficiency anaemia persist with little improvement in recent decades;<sup>4</sup> thinness and food insecurity are still highly prevalent in some contexts, while this generation is simultaneously facing a rapidly escalating crisis of overweight and obesity.<sup>1</sup>

In this paper, we illustrate how a nutrition-focused strategy based on adolescent interventions offers a new lens on tackling the global syndemic of obesity, undernutrition, and climate change,<sup>5</sup> increasingly recognised as an existential challenge to human and planetary health. The conceptual framework for this Series<sup>4</sup> illustrates the diverse influences on adolescents'

nutrition choices, including social influences from peers and family members, economic factors, environmental sustainability, and values of fairness and human rights. Their perspectives, engagement, and creative solutions are likely to be crucial in meeting this challenge. As with other areas of adolescent health, the most effective

## Key messages

- Despite micronutrient deficiencies and food insecurity persisting in many places, and overweight and obesity rapidly increasing, adolescents have been largely overlooked in global nutritional policy frameworks
- Adolescent nutritional problems remain invisible as there are neither established targets nor standardised data collection systems that would inform action
- Adolescent nutritional action for both boys and girls provides a foundation for a healthy start to life for the next generation
- Although multifaceted and multisectoral actions offer great promise, intervention research has overwhelmingly focused on single micronutrients to date
- Greater government fiscal and policy action to both restrict the availability of highly processed foods and enhance healthy and diverse adolescent diets is urgently needed
- Greater retention in education means that schools can provide healthy food environments, nutritionally sensitive social protection at times of crisis, and the skills, knowledge, and motivation to adopt and sustain healthy diets
- Adolescent nutrition advocacy should occur in partnership with young people and be framed within broader commercial, cultural, and ecological contexts

Published Online  
November 29, 2021  
[https://doi.org/10.1016/S0140-6736\(21\)01593-2](https://doi.org/10.1016/S0140-6736(21)01593-2)

This is the third in a Series of three papers on adolescent nutrition

\*Co-lead authors

†Co-leads of the Series (all authors between the co-lead authors and the Series co-leads are listed in alphabetical order)

Mohn Centre for Children's Health and Wellbeing, Imperial College London, London, UK (D Hargreaves MD(Res)); Emergency Nutrition Network, Oxford, UK (E Mates MSc); International Food Policy Research Institute, New Delhi, India (P Menon PhD, P H Nguyen PhD); International Food Policy Research Institute, Washington, DC, USA (H Alderman PhD); Institute for Global Health, University College London, London, UK (D Devakumar PhD); Departments of Global Health, Epidemiology, and Nutrition, Harvard T.H. Chan School of Public Health, Boston, MA, USA (Prof W Fawzi MBBS); Department of Primary Care & Public Health, School of Public Health, Imperial College London, London, UK (G Greenfield PhD); Institute of Community and Public Health, Birzeit University, Birzeit, West Bank, occupied Palestinian territory (W Hammoudeh PhD); Youth Network for the Lancet Standing Commission on Adolescent Health and Wellbeing, Beijing, China (S He MSc); Institute of Economic Growth, New Delhi, India (A Lahiri MTech); UNICEF India, New Delhi, India (V Sethi PhD); Department of Maternal and Child Health, School of Public Health, Peking University, Beijing, China (Z Liu PhD, Prof H Wang PhD); Global Alliance for Improved Nutrition (GAIN), Geneva,

Switzerland (L M Neufeld PhD);  
Murdoch Children's Research  
Institute, University of  
Melbourne, Melbourne, VIC,  
Australia (Prof G C Patton MD)

Correspondence to:  
Dr Dougal Hargreaves, Mohn  
Centre for Children's Health and  
Wellbeing, Imperial College  
London, London W2 1NY, UK  
d.hargreaves@imperial.ac.uk

See Online for appendix

strategies for adolescent nutrition are likely to be both multifaceted and adapted to local context.

Specifically, in this paper we aim to: (1) summarise current understanding of how effective strategies and interventions have been at improving adolescent growth and nutrition; (2) explore how multifaceted strategies, encompassing different sectors and platforms, can be deployed to reach adolescents with effective actions for healthy eating and optimal growth; and (3) offer recommendations on the essential steps for improving adolescent nutrition that can, in turn, support wider societal and planetary health goals.

### What works in adolescent nutrition?

The most effective health actions for adolescents take into account their rapid emotional development and identity formation and the social forces, such as peer context, that drive adolescent decision making.<sup>6,7</sup> These distinctive features of adolescence explain why interventions that might be effective in primary school age children, such as education and behavioural skills training, have a more limited effect in adolescents.<sup>8</sup> Similarly, effective campaigns to influence other adolescent health behaviours, such as smoking, often focus more on short-term impacts, in contrast to campaigns targeted at adults in which long-term risks are highlighted.<sup>9</sup>

Successful adolescent interventions generally resonate with adolescents' values and social context. In the USA and other modern food environments, autonomy, peer approval, and norms have become increasingly important in driving adolescent food choices. Understanding and harnessing these factors could be central to designing effective nutrition policies and programmes for this age group.<sup>8</sup>

Adolescents' greater capacity and willingness to change their behaviour is reflected in greater sensitivity to environmental influences. Migrants who move before or during adolescence are more likely to adopt harmful eating habits of the host country than those who arrive as adults.<sup>10</sup> They are also more likely to adopt harmful eating habits than age-matched counterparts in the host country, suggesting a double vulnerability.<sup>11</sup> Another illustration of openness to change is that the majority of people who adopt a vegan diet (at least in high-income settings) do so in their late teens or early twenties.<sup>12</sup> With careful planning and monitoring to ensure that all nutritional needs are met, this can be a positive choice with many health and environmental benefits.<sup>13</sup>

Adolescent nutritional interventions necessarily vary in their form and effectiveness by age, sex, income level, and geography (eg, rural vs urban), along with social, cultural, and country context. Some interventions will be adolescent-specific, such as those through schools. Others might be adolescent-inclusive but adapted to meet the different social and developmental context of

adolescents. Interventions will also vary by their geographical context in their focus on undernutrition, micronutrient deficiencies, overweight and obesity, or a combination. Lastly, some interventions have proven effectiveness in more than one nutritional context, while many others have a much more limited evidence base.

In the following section, we review existing evidence on approaches to intervention for adolescent nutrition within four broad domains: education and health sectors, the food system, and wider community and social influences (see the table for a summary of key interventions and how they align with the conceptual framework of this Series, and appendix pp 2–3 for search terms and other details of the systematic review of reviews).

### Education sector

Schools provide a platform for a potentially wide range of interventions. The most common and long-standing intervention is school meals (through provision or subsidised sale of food), which have been implemented in diverse ways in both high-income countries and low-income and middle-income countries (LMICs)<sup>14</sup> and have potential for long-term, multisectoral benefits.<sup>5,3</sup> Historically, many such programmes did not include specific nutrition objectives, often included poor-quality foods (both nutritionally and in their appeal to young people), and were not coupled with nutrition education. Reviews suggest such programmes typically had minimal effect on dietary change or nutrition outcomes.<sup>16</sup> Furthermore, many school food programmes focus on younger children and rarely have impact evaluations disaggregated by age group. As a result, their ability to meet the distinct nutritional needs of adolescents (described by Norris and colleagues in this Series<sup>4</sup>) is often not clear.

Over the last decade, several countries have developed and tested new approaches to school food programmes—for example, through policy directives (eg, banning of sugar-sweetened beverages, as in Mexico<sup>26</sup>), menu modifications (eg, improved foods or inclusion of fortified foods, as in Brazil;<sup>24</sup> see also Neufeld and colleagues in this Series<sup>3</sup>), and choice architecture (eg, improving selection, appeal, and prominence of more nutritious food choices, as in the USA).<sup>27–29</sup> The school curriculum offers opportunities to improve dietary diversity and the wider food environment, as well as practical skills around growing and preparing food, skills that are in decline in many food environments.<sup>19,20</sup> In some settings, initiatives such as school gardens and engagement with local food producers can also be valuable, although evidence suggests that impact depends largely on appropriate education linked to the gardens and training for teachers or other school staff to manage them.<sup>18</sup> Aligned with these innovations there is an increasing recognition of the need for a “whole school approach” and promoting nutrition through all available

	Interventions to improve the food and nutrition environment	Interventions to increase nutritional agency*
<b>Education</b>		
Undernutrition	Provision of meals and micronutrient supplements; <sup>14,15</sup> provision of school meals; <sup>15,16</sup> school platforms used to support nutrition of out-of-school adolescents; <sup>17</sup> deworming children and women of reproductive age <sup>15</sup>	Nutrition education (eg, school gardens); <sup>18</sup> knowledge of dietary diversity, food environment, and practical skills; <sup>19,20</sup> use opportunity of school curricula to support nutrition and food preparation <sup>21</sup>
Micronutrient deficiencies	Provision of meals and micronutrient supplements; <sup>14</sup> weekly iron and folic acid supplements to reduce anaemia; <sup>15,22</sup> schools delivering videos (1–2 minutes) in classrooms as motivation for uptake of supplements in clinics; <sup>23</sup> daily iron supplementation (also in malarial areas); <sup>15</sup> provision of micronutrient powders; <sup>15</sup> deworming children and women of reproductive age; <sup>15</sup> inclusion of fortified foods <sup>24</sup>	Nutrition education (eg, school gardens); <sup>18</sup> knowledge of dietary diversity, food environment, and practical skills; <sup>19,20</sup> use opportunity of school curricula to support nutrition and food preparation <sup>21</sup>
Overweight or obesity	Provision of meals and micronutrient supplements; <sup>14</sup> limiting exposure to fast food outlets, as this exposure increases intake of junk food; <sup>25</sup> school platforms used to support nutrition of out-of-school adolescents; <sup>17</sup> decreased consumption of sugar-sweetened beverages, decreased free sugar intake <sup>15,26</sup>	Nutrition education (eg, school gardens); <sup>18</sup> knowledge of dietary diversity, food environment, and practical skills; <sup>19,20</sup> use opportunity of school curricula to support nutrition and food preparation; <sup>21</sup> improving choice architecture <sup>27–29</sup>
<b>Health</b>		
Undernutrition	Nutritional care of HIV-infected children; <sup>15</sup> multiple micronutrient supplementation and macronutrient supplementation for people living with HIV and AIDS; <sup>15</sup> Bright Futures: Nutrition <sup>21</sup>	..
Micronutrient deficiencies	Nutritional care of HIV-infected children; <sup>15</sup> multiple micronutrient supplementation and macronutrient supplementation for people living with HIV and AIDS; <sup>15</sup> Bright Futures: Nutrition <sup>21</sup>	..
Overweight or obesity	Increased potassium and decreased sodium intake to decrease blood pressure <sup>15</sup>	Use of social media to promote healthy behaviour or influence social norms; <sup>30,31</sup> overweight and obesity control interventions <sup>32–34</sup> (including Behaviour Change Communication component, <sup>16,32,35</sup> family component, <sup>36</sup> and longer term or higher intensity interventions <sup>33</sup> )
<b>Food systems, household resources, and regulations</b>		
Undernutrition	Fortification of maize; <sup>15</sup> social protection and cash transfer programmes including nutrition education <sup>25,37</sup>	..
Micronutrient deficiencies	Cash transfers to increase uptake of healthy meals and micronutrient supplementation <sup>38</sup>	..
Overweight or obesity	Regulations to limit access to unhealthy meals, drinks, or snacks during the school day; <sup>39</sup> reducing the impact of marketing of foods and non-alcoholic beverages on children; <sup>15</sup> subsidies and taxes to reduce the intake of unhealthy beverages and foods; <sup>40–43</sup> minimum alcohol pricing and increased taxes on sugar-sweetened beverages <sup>44</sup>	..
<b>Social and community influences</b>		
Undernutrition	..	Increasing community ownership of interventions, inclusive of culturally relevant information; <sup>45</sup> mentoring from community members to increase impact of interventions <sup>46</sup>
Micronutrient deficiencies	..	..
Overweight or obesity	Policies to encourage active transport and safe walking or cycling <sup>47</sup>	Youth-led movements—eg, Bhalo Khabo Balo Thakbo (Eat Well Live Well); Real Food Systems Youth Ambassadors <sup>48</sup>
<b>Multisectoral</b>		
Undernutrition	..	..
Micronutrient deficiencies	..	..
Overweight or obesity	Addressing overweight and obesity through multiple interventions (schools, parent engagement, etc) <sup>49,50</sup> —eg, Joint Strategy to address obesity in China; <sup>51,52</sup> reductions of overweight and obesity through interventions of the Amsterdam Healthy Weight Programme <sup>47</sup>	..
*Adolescent agency was operationalised in the <i>Lancet</i> 2016 Commission <sup>6</sup> across three aspects: (1) having the skills and knowledge to advance the conversation (across the board, young people, and decision makers); (2) having forums available for young people to engage; and (3) ensuring there are resources available for engagement.		

**Table: Mapping evidence-based interventions to the conceptual framework**

means.<sup>54</sup> This includes changes to class curricula to embed nutrition; facilitation of physical activity and active transport; creating a healthy food environment; supporting monitoring, screening, and supplementation programmes (as needed); and reinforcement through individual engagement by school staff.<sup>55,56</sup>

Beyond food, schools provide an important platform for delivery of interventions targeted to high-risk groups, such as weekly iron and folic acid (WIFA) supplementation for adolescent girls. Anaemia remains an important public health challenge in many countries and is associated with increased fatigue, difficulties in learning, higher risk of infections, and an increased risk to both mother and baby in future pregnancies. In populations with a high prevalence of iron deficiency anaemia, WIFA can reduce anaemia by 35% in adolescent and adult women.<sup>22</sup> Such programmes appear cost-effective when compared with economic losses due to iron deficiency anaemia.<sup>23,56</sup> However, WIFA programmes have been implemented in only a handful of countries, anaemia in adolescents boys has been absent from the data and action agenda, and to date, most countries have seen no or only minor reductions in adolescent anaemia.<sup>57</sup> One reason for the lack of progress is that anaemia is multicausal and the proportion of anaemia due to iron deficiency varies substantially by context.<sup>58</sup> There is therefore a need to improve the quality of data tracking anaemia, to ensure that programmes can address its different causes.<sup>59,60</sup>

Although global school enrolment has increased considerably since 2000 and is a central platform for action to improve nutrition, 37% of the secondary school-age population in sub-Saharan Africa never attend school and many of those enrolled are irregular attendees.<sup>61</sup> New approaches are needed to reach out-of-school adolescents and those who rarely attend to improve nutrition literacy, ensure food and nutrition security, and address specific nutrition issues. Ghana and India offer examples of WIFA programmes in which both adolescents in school and those who are out of school are included. In India, the Anganwadi centres offer networks of peer support groups and women's self-help groups.<sup>17</sup> Elsewhere, schools have delivered videos of 1–2 minutes in classrooms to encourage the take-up of supplements at clinics, thus having the potential to reach parents and siblings as well as neighbours of school pupils.<sup>23</sup> Take-home rations from school feeding programmes have also been used to reach other vulnerable household members.

The contribution of inequitable gender norms and associated high levels of early marriage and teenage pregnancy to poor nutrition in adolescent girls and their offspring requires concerted action. Continued engagement with education is arguably the most effective nutritional intervention available in some low-resource settings. In addition, school curricula offer an underused opportunity for acquiring skills in childcare, nutrition, and food preparation.

### Health sector

Health professionals can be a trusted source of nutrition advice for adolescents, providing information and implementing screening and treatment programmes

that can be adapted both to the individual's needs and to the local setting. Interventions in health-care settings, or those delivered by health professionals in community or education settings, are therefore an important component in adolescent nutrition strategies.<sup>21</sup> In more severe cases, undernutrition or overweight and obesity might be a primary reason for seeking health care; in other cases, nutrition, eating, and growth problems might be identified as part of a screening programme, or opportunistically when a clinician is assessing another problem. Aside from WHO's Global Accelerated Action for the Health of Adolescents (AA-HA!) guidance,<sup>62</sup> there are few clinical guidelines targeted towards adolescents; depending on the setting, the use of nutritional supplements and interventions such as treatment for parasites might be guided by population-based screening programmes, clinical assessment, or laboratory investigations. In all settings, an opportunistic approach to nutrition assessment and counselling during all contacts with adolescents is key. Specialist interventions in the treatment of eating disorders or medical complications of obesity, including gastric banding, are important for a small proportion of adolescents but are beyond the scope of this paper.

A wide variety of individual and group-based interventions have been developed for overweight and obese adolescents, including: nutrition advice and supervised diet programmes; behavioural approaches such as motivational interviewing;<sup>32</sup> physical activity interventions; and programmes which combine some or all of these components.<sup>33,34</sup> A review of 18 studies with patients aged 4–18 years, mostly in the USA and other high-income countries, found that primary care interventions led to a small but significant reduction in body-mass index (BMI; 2–11% reduction in BMI Z-score). Interventions appeared to be more effective when they included a behaviour change or psychology component<sup>16,35</sup> or a family component.<sup>36</sup> Longer duration and higher intensity interventions were also more likely to be effective.<sup>31</sup> In the smaller subset of adolescent studies, the effects of primary care interventions on obesity were mixed; the only study with a clear impact involved an intensive programme of 28 contacts over a 5-month period.<sup>63</sup>

Neufeld and colleagues<sup>3</sup> highlight the growing influence of social media on dietary choices, body image, and psychological wellbeing, both through advertising and marketing to adolescents and subsequent peer interactions. These same processes underpin interest in e-health and m-health interventions for overweight and obese adolescents, although most are management-focused rather than prevention-focused. This is a rapidly changing field in which interventions range from those intended to increase knowledge, through to self-monitoring apps providing immediate feedback on performance, which harness the reach of social media to promote healthy behaviour and influence societal

norms.<sup>30,31</sup> Proponents highlight young people as early adopters of new technology and the potential of social media to shape peer norms in a life phase during which peer influence is greater than at any other point in the lifecycle. The potential has grown, with mobile phones and computers used in almost every area of young people's lives in both high-income countries and many LMICs. Although digital health interventions for adolescent nutrition are promising areas of research, consistent impact has yet to be demonstrated.

Health interventions combining dietary, physical activity, and behavioural strategies with family and perhaps digital components have potential to reduce overweight and obesity in adolescents.<sup>36</sup> However, wide variation in the effect of apparently similar interventions suggests that further work is needed to refine and improve these interventions in a way that addresses the needs of different adolescent groups. For example, physical activity organised or delivered by peers appears to bring greater engagement, sense of responsibility, and retention among adolescents in disadvantaged communities than adult-led activities.<sup>64</sup> Similarly, one of the issues highlighted in a few studies among First Nations, Inuit, and Métis communities in Canada is that interventions are more effective if they incorporate culturally relevant nutrition information and if the community has a sense of ownership.<sup>45</sup>

### Food systems, household resources, regulations, and food environments

A recurring limitation of nutrition interventions in the health and education sectors is that the benefits of increased information or motivation can be outweighed by additional costs or practical barriers to making healthy choices.<sup>65</sup> Additionally, these limitations can sometimes be exacerbated by commercial exploitation of adolescents as a potentially vulnerable group, particularly through social media.<sup>3</sup> Supply-side interventions, such as farming subsidies and other agriculture interventions, can have large influences on food availability and consumption at population level and illustrate the key role of governments in balancing commercial and public health interests.<sup>66,67</sup> Similar potential impact and political tensions apply to food regulation—for example, WHO's campaign to ban trans-fatty acids,<sup>68</sup> initiatives to regulate front-of-pack nutrition labelling,<sup>69</sup> and targeted taxes and policies to reduce the intake of sugar-sweetened beverages are all important initiatives with potential for substantial impact at population level.

A meta-analysis based largely on intervention studies supports the efficacy of subsidies to increase consumption of healthy foods and taxation to reduce intake of unhealthy beverages and foods in adults and children. A 10% decrease in price increased consumption of healthy foods by 12%, whereas a 10% increase in price (ie, tax) decreased consumption of unhealthy foods by 6%.<sup>70</sup> For example, in Seattle, USA, sales volumes of

sugar-sweetened beverages fell by 22% following introduction of a new tax;<sup>40</sup> in the UK a new tax was effective in reducing the sugar content of available drinks and total sugar consumption;<sup>41,42</sup> and Chile's introduction of new regulations in 2014 was associated with a 23.7% reduction in purchase volume of beverages that were high in sugar, sodium, or fats.<sup>43</sup> A combination of taxation, regulation, and public education campaigns appears to maximise impact.<sup>71,72</sup>

There is sparse evidence on the specific impact of these nutrition interventions on adolescents. Screen advertising bans appear to have more consistent impact on primary school age children,<sup>73,74</sup> while there is some evidence that restricting tobacco and alcohol advertising might have greater impact on adolescents than adults.<sup>75,76</sup> Evidence from minimum alcohol pricing and additional taxes on sugar-sweetened beverages shows disproportionately large impact among those with lower disposable income, reflecting a greater sensitivity to price, and among those with less established patterns of consumption;<sup>44</sup> both would favour greater impact among young people than older adults.

Equally important are demand-side interventions to improve people's ability and motivation to buy nutritious food. In both high-income countries and LMICs, economically vulnerable families struggle to provide healthy diets for children in their homes; indeed, child hunger and poor diets at home are key challenges worldwide. The COVID-19 pandemic and associated lockdowns have exacerbated pre-existing food insecurity in many countries—often disproportionately affecting young people who have less secure employment and fewer resources to fall back on. US data show that food insecurity in early adolescence predicts more rapid increase in BMI by the age of 31 years;<sup>77</sup> in addition to health impact, it can also reduce educational engagement and attainment.<sup>78</sup> In rural Tanzania, food insecurity among adolescents was linked to illness and ability to take on adult farming responsibilities (which can involve greater energy expenditure)<sup>79</sup>—a cycle that could be exacerbated as the increasing effects of climate change are seen.<sup>80</sup>

Social safety net programmes play a key role in supporting families around food and nutrition and in ensuring that children can access schools and complete their education. Many of these social transfer programmes, such as the food assistance programmes in the USA that include nutrition education for families with children, have been shown to be effective in improving family and child diets.<sup>25,37</sup> Globally, economically developing and nutritionally transitioning countries spend an average of 1.5% of gross domestic product on these programmes. With some regional variation, on average, spending on safety nets increases as a share of gross national product with national income. Cash transfers (which can be conditional or unconditional) represent an increasing share of

transfers in most regions<sup>81</sup> but globally only a few cash transfer programmes include adolescents in the target group, such as South Africa's Child Support Grant.<sup>82</sup> Even without being explicitly targeted towards adolescents, there is good evidence that cash transfers can be effective in influencing wider adolescent outcomes, including extending schooling, delaying parenthood, and reducing HIV risk.<sup>38</sup> Emerging evidence suggests that cash transfers could also be a powerful tool to support adolescent nutrition in some contexts, with potential to combine healthy meals and micronutrient supplementation with the wider educational and social goals listed previously, but current evidence is limited and a potential caveat is that cash transfers can lead to excess weight gain.<sup>38</sup> As with interventions in other sectors, cash transfers clearly need to be planned as part of an integrated strategy that is designed and evaluated according to local needs and priorities. Additionally, developmental stages must be considered, taking into account how different priorities for spending change through early, mid, and late adolescence.<sup>8</sup>

Even where healthy food is available and affordable, adolescent choices can be heavily influenced by commercial influences. These include advertising and marketing factors (eg, clarity of labelling, supermarket placement, and social media influences) as well as convenience. Neufeld and colleagues<sup>3</sup> describe how the proximity to schools of shops selling unhealthy food can influence adolescent food choices and result in higher consumption patterns.

### Social and community influences

The social brain of adolescents has been well described<sup>83</sup> and food choices illustrate the different levels at which these social influences can operate, from peer groups and local communities, to virtual networks and international social movements.

Community-grounded interventions offer opportunities to work with a wide range of local institutions. Community organisations can both deliver interventions and support strategies that improve local food environments and food choices, from non-governmental organisations and religious communities, to local businesses and sports or music clubs. Many of the most effective obesity interventions are designed to be delivered in community settings, taking into account a specific social context.<sup>35</sup> For example, mentoring from members of the African American community increased the impact of nutrition interventions for African American young people in one US study.<sup>46</sup> Unfortunately, despite their enormous potential, there is a relatively sparse literature evaluating the impact of interventions based in these community sector organisations, to date.

Community interventions provide opportunities for targeting social norms and the environmental context in which nutritional choices are made. Firstly, a range of

policies to promote active transport, including cycle lanes, pedestrianised zones, and safety measures for those walking or cycling, have been implemented by national and local governments and are often targeted specifically at young people.<sup>47</sup> Secondly, local governments can influence planning rules, such as restrictions to fast-food restaurants near schools or licensing of street-food vendors. For example, exposure to fast-food outlets was strongly associated with junk food intake in one study of Canadian adolescents.<sup>84</sup>

Looking at social influences more broadly, adolescents today have opportunities to link to a much wider range of social networks than previous generations. Information technology and social media have transformed the way in which both social and commercial influences in adolescence are expressed and experienced. Although social media has sometimes been linked to harmful nutrition outcomes—for example, in reinforcing weight related-stigma and normalising unhealthy body images—it has also opened up opportunities for young people to connect with people outside their immediate community and has transformed social opportunities for those with limited mobility.

Social media illustrates a need to shift from viewing adolescents as passive recipients of interventions to active partners pursuing shared goals. The work of the Real Food Systems Youth Ambassadors<sup>48</sup> shows how young people are playing increasing leadership roles in new social movements, which often combine local action with networking and campaigning through social media. Similarly, Bhalo Khabo Bhalo Thakbo (Eat Well Live Well) is a social media campaign led by adolescents and supported by over 5 million people in Bangladesh. It targets both individual responsibility, whereby supporters take a pledge to buy and eat more healthy food, and a system-level goal to “change the way food is produced, manufactured and sold”; both political and consumer power of young people is explicitly mobilised to campaign for change.

### Multifaceted interventions to address multiple sectoral challenges to better nutrition

Although the literature reviewed in the previous section presents some evidence for the effectiveness of specific nutrition interventions (summarised in the table and the WHO e-Library of Evidence for Nutrition Actions [eLENA]),<sup>15</sup> there is a compelling case that no single intervention or platform will be effective alone. Just as the importance of integrated interventions to improve maternal and child nutrition is increasingly recognised,<sup>85</sup> the forces shaping adolescent growth and dietary intake demand coordinated multifaceted action across sectors in order to achieve the greatest benefits. Furthermore, the multifaceted action at local and national levels must reflect the diverse contexts in which adolescents live.<sup>3</sup> Norris and colleagues<sup>1</sup> highlight the limited progress in achieving multiple nutrition goals: no country has been

For Bhalo Khabo Bhalo Thakbo  
see <https://www.bhalokhabobhalothakbo.com/>

successful in preventing an unhealthy nutrition transition in which one set of problems (underweight) is replaced by another set—ie, overweight and obesity, often with ongoing concerns about micronutrient deficiencies and inequalities. To date, no country has managed to either reverse the trend or achieve an ideal transition. There are several reasons for this: a lack of timely adaptation of policy settings to a different nutritional reality; a failure to adopt double-duty or triple-duty actions; the failure to target this age group before metabolic adaptations and obesity become entrenched; and most importantly, the piecemeal and fragmentary approach in which simple, single-faceted interventions have generally predominated.

Emerging evidence of the potential impact of a multifaceted approach is illustrated by two strategies to reduce overweight and obesity among children and young people in China and the Netherlands. In China, a team at Peking University developed a comprehensive intervention to promote knowledge, healthier diet, and physical activity (drawing on literature reviews<sup>51,86</sup> and engagement work with local stakeholders). Following mixed results from an initial cluster randomised study,<sup>52</sup> a further trial was designed with greater emphasis on engaging teachers, families, and members of the local community to promote adherence.<sup>87</sup> From baseline to 9 months, the mean BMI decreased in the intervention group, whereas it increased in the control group (mean difference  $-0.46$  kg/m<sup>2</sup> [95% CI  $-0.67$  to  $-0.25$ ;  $p < 0.001$ ]). The intervention was also beneficial for other adiposity outcomes, dietary and sedentary behaviours, obesity-related knowledge, and physical fitness (all  $p$  values  $< 0.05$ ). A larger effect was seen on dietary behaviours than on physical activity. Subsequently, local education bureaux have incorporated the intervention into regular school health services and further refinements are now underway, with a view to addressing community and local environmental factors such as access to convenience stores.<sup>88</sup>

Another example of a multifaceted, multisectoral intervention is the Amsterdam Healthy Weight Programme, which was introduced in 2013 in response to local concerns about the prevalence of childhood overweight and obesity. The programme took a whole systems approach with a wide range of complementary interventions designed to promote healthier diets, physical activity, and good quality sleep—all based on a philosophy that “the healthy choice should be the easy choice” for children and young people, parents, and everyone in the community.<sup>47</sup> Early data showed that obesity prevalence in the most deprived group of 0–18 year olds fell by around a quarter between 2012 and 2015 (8% to 6%), while the total prevalence of overweight and obesity fell from 21% to 18.5%.<sup>47</sup> Although the absolute reduction in obesity is relatively small, these findings should be seen in the context of rising trends in many comparable cities and the

difficulty in achieving rapid change when obesity has been rising steadily over decades. However, longer-term data on the sustainability and impact of this approach are needed, along with lessons for other cities and information on specific impact for adolescents and younger children.<sup>89</sup>

Similar multifaceted local strategies have also reported some success in reducing childhood obesity (0–5 years) in Leeds, UK,<sup>49</sup> and in four US projects. In the USA, strategies in New York City and Philadelphia targeted students in grades K–8 (ie, age 5–14 years). In New York City, obesity in this age group over the period 2006–07 to 2010–11 declined from 21.9% to 20.7%, representing a relative decline of 5.5% ( $p < 0.001$ ). In Philadelphia, from 2006–07 to 2011–12, obesity declined from 21.5% to 20.5%, representing a 4.7% relative decrease (7.7% for severe obesity;  $p < 0.001$ ).<sup>50</sup> Emerging efforts in India, another country with a large population of adolescents and young adults, that were developed using insights from a nationwide survey are described in panel 1.

### Data-informed governance for intersectoral action

Standardised and more comprehensive data are key to effective intersectoral action and accountability of policy makers. Too often, adolescent nutritional problems remain invisible, as there are neither established targets nor standardised data collection systems that would inform action (figure and appendix pp 4–5). Data should also inform the design and delivery of multifaceted interventions for adolescents, which reach them at the right time, in the right place, with the right interventions for their life stage. Similarly, data allow targeted responses to the specific nutritional needs of different populations, supporting a shared understanding and a shared commitment across sectors, coupled with focused actions within sectors to resolve key bottlenecks.<sup>93–95</sup> When multifaceted, multisectoral solutions reach adolescents where they live, study, play, and eat, they have great potential. Therefore, data-informed governance for actions across sectors must hold the adolescent at the centre of all planning and must align actions in ways that address the drivers of poor nutrition for adolescents at each stage of their life and within each sector.<sup>6</sup>

The use of data and local insight to drive intersectoral action should start with strategy-focused diagnostics: what are adolescents eating; what is their nutritional status; what is happening in their schools; what are families with adolescents struggling with; what do local food retail environments look like; what is happening within the health sector; and how are national policies taken up differently in different places? Adolescent nutrition profiles will differ between different food environments, reflecting variations in available food, social expectations, and the degree of autonomy exercised by adolescents in different settings.<sup>3</sup> Second, with greater

**Panel 1: Use of data to generate multifaceted policy actions for adolescent nutrition in India**

India is home to 253 million adolescents and has the largest number of adolescents in any country, globally. Despite recognition that adolescent nutrition is important, there has long been an absence of data on (1) specific age groups (10–14 years); (2) specific outcomes and risk factors, such as micronutrient deficiencies, worm infestation, non-communicable disease risk, anthropometry, physical activity, mental health, and household food security;<sup>30</sup> and (3) school-based services. Another data challenge was limited use of gold standard methodologies.

The Comprehensive National Nutrition Survey (CNNS)<sup>31</sup> 2016–18 addressed these gaps head on. It used robust tools and methods to provide nationally representative data on the nutrition status of preschoolers (0–4 years), school-age children (5–9 years), and adolescents (10–19 years). Surveying 35 856 adolescents aged 10–19 years, and a subset of 16 181 adolescents for biological samples, the CNNS revealed numerous adolescent nutrition challenges, including underweight, anaemia, overweight, poor diet quality, and low physical activity (appendix pp 6–8). It also highlighted limited coverage of school-based services and inequities among rural and poor populations. The findings clearly called for multifaceted actions cutting across food systems, health, and education sectors.

Building both on these findings and on a series of policy engagements, the Government of India is gearing up to strengthen nutrition components in ongoing programmes reaching adolescent boys and girls, and to enact legislation to ban unhealthy foods in the areas within and around school campuses.

- Multi-stakeholder policy and research groups will further analyse the data to answer policy questions on coverage, risk factors, and determinants of anaemia and poor adolescent nutrition.
- Data from the report are being used to back health and promote peer-to-peer activities under the Eat Right School initiative of the Food Safety Standards Authority of India, to create an environment of safe food and healthy eating for students and teachers in various schools.<sup>32</sup> These data are also being used to back legislation under development aiming to regulate the marketing of unhealthy foods in schools.
- A new school health curriculum has been launched to ensure that 1 hour per week and 26 hours in a year are devoted to health and nutrition activity-based learning, across government schools.
- Health and nutrition promotion activities in schools, covering physical activity; reducing accessibility of unhealthy foods in schools; mass campaigns around healthy snacking; and reducing salt, sugar, sweets, and fried foods are gaining momentum under the Eat Right India movement.

Overall, the engagement around the CNNS highlights the value of investing in and closely examining data, conducting relevant analysis, and directly engaging with policy stakeholders across sectors. This has allowed an acknowledgment of the challenge of nourishing India's adolescents and identification of specific policy actions for the short term and long term.

shared understanding, barriers and actions within each sector or area could be identified and acted upon. For example, if school meals are poor because national nutrition guidelines are suboptimal, then guidelines will need revision; if families with adolescents are struggling financially, then social protection measures might be needed; if adolescents are encountering only unhealthy foods in their food environment, then market-focused efforts will be needed. Third, regular monitoring of locally grounded and adolescent-centred action plans are essential. To build local and national accountability, it will be crucial to engage adolescents, their families, their schools, and local government to review and identify priorities, set targets, and track progress.

The major available global databases on adolescents around the world include those maintained by agencies such as UNICEF and WHO, as well as those hosted and maintained by national governments mainly in high-income countries (see figure and panel 1). More disaggregated data on adolescents and greater standardisation of data between countries would support a more data-driven approach to adolescent nutrition policies. Of the domains in the framework for adolescent

nutrition in our Series, data are most sparse around economic development, urbanisation, and food and agriculture systems, and how they intersect with adolescents. Similarly, the key outcomes we emphasise in this Series—diets and nutritional effects and wellbeing—are also priorities for investment in better data. Creativity is needed to construct databases that consolidate data from diverse sources in ways that help us understand how the domains of the framework play out for adolescents and young adults. Panel 1 and appendix pp 4–5 illustrate how a recent investment in such a multisectoral data strategy in India has helped to identify priority actions and inform multifaceted local and national strategies.

**Recommendations**

The world's adolescents are at a nutritional tipping point. Long-standing problems of micronutrient deficiencies, low weight and food insecurity, and short stature persist, as newer problems of obesity and metabolic disturbances rapidly emerge. Pervasive neglect in policy, programming, and research means that the field lacks knowledge, systems, and human resources compared with other age



	Population Council Adolescent Data Hub	Demographic and Health Survey (DHS) data	Health Behavior in School Age Children (HBSC)	United States Adolescent Health	WHO Maternal, Newborn, Child and Adolescent Health data portal	UNICEF adolescent health data (Multiple Indicator Cluster Survey)
<b>Databases</b>	<ul style="list-style-type: none"> <li>• 127 LMICs</li> <li>• 473 datasets (453 observational, 20 experimental, 358 national representative data)</li> <li>• Age 10–14, 15–19, and 20–24 years</li> </ul>	<ul style="list-style-type: none"> <li>• All DHS surveys conducted from 2000 to 2017 with availability of data for girls or boys age 15–19 years</li> </ul>	<ul style="list-style-type: none"> <li>• WHO collaborative cross-sectional national survey</li> <li>• Data was collected every 4 years on 11–15-year-olds in 49 countries and regions across Europe and North America</li> </ul>	<ul style="list-style-type: none"> <li>• US national and state level</li> <li>• Age 9–20 years</li> </ul>	<ul style="list-style-type: none"> <li>• Global health data, including regional and country data</li> <li>• Age 10–19 years</li> </ul>	<ul style="list-style-type: none"> <li>• More than 100 countries around the world</li> <li>• Age 15–19 years</li> </ul>
<b>Domains</b>						
Natural systems and planetary health	No data	No data	No data	No data	No data	No data
Social, cultural, and gender norms	<ul style="list-style-type: none"> <li>• Gender attitudes and beliefs</li> <li>• Community engagement</li> <li>• Crime</li> <li>• Social networks</li> <li>• Subjective expectations</li> </ul>	No data	<ul style="list-style-type: none"> <li>• Social environments</li> </ul>	<ul style="list-style-type: none"> <li>• Healthy relationships</li> </ul>	<ul style="list-style-type: none"> <li>• National policies</li> </ul>	No data
Economic development, urbanisation, and food and agriculture systems	<ul style="list-style-type: none"> <li>• Economics</li> <li>• Health-care access and utilisation</li> <li>• Migration and mobility</li> </ul>	<ul style="list-style-type: none"> <li>• Iron supplementation</li> </ul>	No data	No data	No data	No data
Household, school, and peer dietary and activity patterns	<ul style="list-style-type: none"> <li>• Demographic characteristics</li> <li>• Education</li> <li>• Family and household structure</li> <li>• Time use</li> </ul>	No data	No data	<ul style="list-style-type: none"> <li>• Demographics</li> </ul>	No data	<ul style="list-style-type: none"> <li>• Education</li> <li>• Marriage</li> <li>• Early demographics</li> <li>• Participation</li> </ul>
<b>Individuals</b>	Physical activity	No data	No data	No data	No data	No data
	Cognitive and emotional growth	<ul style="list-style-type: none"> <li>• Mental health</li> </ul>	No data	No data	<ul style="list-style-type: none"> <li>• Mental health</li> </ul>	<ul style="list-style-type: none"> <li>• Mental health</li> </ul>
	Food choice and dietary intake	No data	<ul style="list-style-type: none"> <li>• Women's diet diversity score</li> </ul>	No data	No data	No data
<b>Other relevant outcomes</b>	<ul style="list-style-type: none"> <li>• Physical health</li> <li>• Reproductive health</li> </ul>	<ul style="list-style-type: none"> <li>• Anaemia</li> <li>• Anthropometry</li> <li>• Adolescent childbearing</li> </ul>	<ul style="list-style-type: none"> <li>• Health and wellbeing</li> <li>• Health behaviours</li> </ul>	<ul style="list-style-type: none"> <li>• Reproductive health</li> <li>• Physical health and nutrition</li> <li>• Substance abuse</li> </ul>	<ul style="list-style-type: none"> <li>• Adolescent sexual and reproductive health</li> <li>• Morbidity</li> <li>• Mortality and cause of death</li> </ul>	<ul style="list-style-type: none"> <li>• Childbearing</li> <li>• Violence</li> </ul>

Figure: A review of global databases on adolescent health

LMICs=low-income and middle-income countries.

groups (see panel 2 for research recommendations). More than ever before, adolescent nutrition is tied to global changes in urbanisation, food systems, industry marketing, and digital technologies, accompanied by planetary change and more recently pandemic disruptions. We make the following recommendations for action to ensure adolescents' own health, and that of the next generation they will be caring for, while minimising environmental impacts.

### Build commitment to adolescent nutrition through evidence-based and accountable systems

#### 1. Targets should be established for adolescent nutrition in its global tracking and accountability mechanisms

Optimal nutrition of adolescent girls and boys is a pillar of human capital, but no global targets have been set for either undernutrition or overweight and obesity. The research community, UN agencies, international non-governmental organisations, donors, and national

**Panel 2: Research needs to support better actions for adolescent nutrition**

We base several of the recommendations in this Series on an evidence base that is still too limited. Too much remains unknown, especially around specific platforms and specific age groups. Some critically important areas for further investment are described below, focusing on neglected age groups, neglected intervention areas, and neglected platforms:

- Age groups: data and research on nutrition challenges and determinants appears to be most lacking for males and for the 10–14 years age group, even though, programmatically, many countries have school platform-based programmes reaching this age group. Programmatically, the later adolescent age groups are most neglected. Programmes for secondary school students will reach the middle group of 15–19 year olds, but both data and programmes for young adults 20–24 years of age are largely focused on reproductive health and childbearing. All age groups are aggressively targeted by commercial marketing, despite this neglect in academic research and government programmes.
- Intervention areas: across the range of intervention areas, school interventions are probably the best researched, but there remain many questions about health platforms and adolescents, including on how to tailor antenatal care services for adolescents in ways that enable access, engagement, and inclusivity. Similarly, there are large gaps in our understanding of the full range of effects of multifaceted interventions such as those tested in China and recently launched in India. In large countries such as these, tailored and context-specific research efforts will also be important to inform policy and programme actions.
- Platforms: reaching out-of-school adolescents more effectively is a critical need, and current research is scarce. Depending on the country, alternative platforms to reach this group could include community groups and non-governmental organisations, juvenile justice systems, refugee services, and homeless health services. Similarly, little is known about platforms to improve nutrition and food choices of adolescent boys, as the focus is almost always on girls and young women. Additionally, although social media is not a neglected platform for research in general, it has not been harnessed enough around topics such as nutrition; this is a major gap given the high and expanding use of social media among young people around the world.

Across all these areas, research will need to be context-specific, ensuring adequate attention both to the range of food environments and to the range of school and health environments that adolescents are reached by.

governments will only bring about the radical changes needed if data-based accountability mechanisms are established.

*2. National data collection systems should be strengthened and extended to include standardised indicators and determinants of adolescent nutritional status (eg, dietary intake), and coverage of interventions among both adolescent boys and girls*

Most countries lack data to monitor adolescent nutrition and ensure timely action in the context of rapid global shifts. Data should be central to national efforts to prioritise action and to assess progress in adolescent nutrition and development.

*3. Donors and research funding agencies should invest in and prioritise existing knowledge gaps to accelerate action for adolescent nutrition*

There are glaring gaps in our understanding of adolescent nutrition, ranging from its consequences for maturation

of physiological systems through to the shifting determinants of adolescent food choices and how to move action beyond single micronutrient supplements (which do not address the underlying causes of nutritional deficiencies), to the multiple drivers of food choice.

**Enhance policy and programmatic actions that favour healthy adolescent nutrition**

*4. National governments and relevant state jurisdictions should use fiscal and policy levers to promote healthy diets for children and adolescents*

These levers can be used to increase the availability and affordability of healthy foods; they can also be used to restrict the marketing of, reduce access to, and extend taxation of increasingly ubiquitous inexpensive unhealthy foods.

*5. Researchers should partner with consumer and shareholder activists to hold food manufacturers and distributors, from global to local, accountable for their food policies and marketing directed to children and adolescents*

Across all countries, adolescents are increasingly consuming unhealthy foods that are high in sugar, fats, and salt, and of low nutritional value. All actors in the food system must support healthy food environments and restrict sales of unhealthy alternatives.

*6. National and jurisdictional governments should guarantee the availability of healthy food through school meals, particularly in settings of poverty and food insecurity, and at times of crisis*

Many adolescents and their families live in extreme poverty, and climate environmental changes are likely to bring even greater food insecurity. The expansion of education and the integration of nutrition in educational settings offers the opportunity to extend nutritionally sensitive social protection schemes.

*7. Ministries of education should mandate the creation of healthy school food environments, including the enforcing of nutritional standards for school meals; subsidising of nutritious foods; restricting the sale and advertising of processed foods; and encouraging engagement with the local community, including parents and farmers*

Nutrition is central for school learning and ultimately human capital. Equally, schools can play an important role in adolescents' nutrition through the curriculum, the foods consumed at school, and the nutritional norms adopted by the student group.

*8. Ministries of education and schools should expand school curricula, for both girls and boys, in ways that extend to relevant skills-based training, including understanding the production of nutritious and sustainable foods, and the scope for adapting local agriculture to environmental change*

Particularly where aligned with the interests of youth in rural and farming communities, to support them as

champions of change for nutrition, school curricula offer opportunities beyond the promotion of developmentally appropriate food literacy for both boys and girls, to skills-based training in maintaining traditional knowledge and renewing skills around preparation and cooking of nutritious and delicious food.

**9. National and subnational governments should ensure young people have easy access to accurate nutrition information through simple, bold, front-of-pack warning labels, as well as promoting healthy diets on social media with messages grounded in an understanding of the drivers of adolescent food choice**

Increasingly, food choices are shaped by mass and social media, amplified by food, beverage, and fashion industries, which shift norms around sexual attractiveness, body image, and physical and mental health.

#### Place adolescent nutrition advocacy within a broader ecological context

**10. All nutrition constituents should partner with young people in advocacy for diets that are healthy, safe, and sustainable**  
Highly processed foods are bad for people and the planet: reducing consumption will be beneficial for both human and planetary health. Advocacy for healthy adolescent nutrition needs to address commercial, structural, and ecological drivers of food choice, and use the agency of young people to bring about transformative change.

#### Contributors

DH, EM, and PM contributed equally to the manuscript. DH, PM, and GCP conceptualised the manuscript; DH, EM, PM, LMN, and GCP did the final manuscript preparation. DD, EM, GG, WH, DH, HA, and WF developed the section on the evidence base of interventions. PHN, VS, AL, and PM developed the India case study; ZL and HW developed the China case study. PHN and PM reviewed availability of global data. SH contributed the youth perspective. All authors drafted sections of the manuscript, contributed to the interpretation and discussion, and approved the final manuscript.

#### Declaration of interests

DH was the Deputy Chief Scientific Adviser at the Department for Education (England) from September, 2020, to September, 2021. HA reports grants from the Bill & Melinda Gates Foundation, outside the submitted work. WH reports grants from Fondation Botnar, outside the submitted work. All other authors declare no competing interests.

#### Acknowledgments

This work received funding support from Fondation Botnar and the Wellcome Trust. Neither organisation played any role in writing the manuscript or the decision to submit for publication. We would also like to acknowledge support from the following: the National Institute for Health Research (NIHR) through the National School for Public Health Research Programme and the Applied Research Collaboration North West London (DH); the Bill & Melinda Gates Foundation (PHN, PM, HA); the National Key R&D Programme of China (2016YFC1300204) and the National Natural Science Foundation of China (81903343; ZL, HW); and Ireland's Department for Foreign Affairs (EM). The Comprehensive National Nutrition Survey (CNNS) 2016–18 was conducted under the leadership of the Ministry of Health and Family Welfare (MoHFW), Government of India and Technical Advisory Committee designated by the MoHFW, in collaboration with UNICEF and the Population Council. The CNNS was funded through a generous grant from Aditya Mittal, president of ArcelorMittal, and Megha Mittal, Managing Director of ESCADA. The views expressed in this publication are those of the

authors and not necessarily those of any funding bodies or institutions mentioned above.

Editorial note: the *Lancet* Group takes a neutral position with respect to territorial claims in published maps and institutional affiliations.

#### References

- Norris SA, Frongillo EA, Black MM, et al. Nutrition in adolescent growth and development. *Lancet* 2021; published online Nov 29. [https://doi.org/10.1016/S0140-6736\(21\)01590-7](https://doi.org/10.1016/S0140-6736(21)01590-7).
- Drake LJ, Lazrak N, Fernandes M, et al. Establishing global school feeding program targets: how many poor children globally should be prioritized, and what would be the cost of implementation? *Front Public Health* 2020; **8**: 530176.
- Norris SA, Frongillo EA, Black MM, et al. Nutrition in adolescent growth and development. *Lancet* 2021; published online Nov 29. [https://doi.org/10.1016/S0140-6736\(21\)01590-7](https://doi.org/10.1016/S0140-6736(21)01590-7).
- Neufeld LM, Andrade EB, Suleiman AB, et al. Food choice in transition: adolescent autonomy, agency, and the food environment. *Lancet* 2021; published online Nov 29. [https://doi.org/10.1016/S0140-6736\(21\)01687-1](https://doi.org/10.1016/S0140-6736(21)01687-1).
- Swinburn BA, Kraak VI, Allender S, et al. The Global Syndemic of Obesity, Undernutrition, and Climate Change: The *Lancet* Commission report. *Lancet* 2019; **393**: 791–846.
- Patton GC, Sawyer SM, Santelli JS, et al. Our future: a *Lancet* commission on adolescent health and wellbeing. *Lancet* 2016; **387**: 2423–78.
- Dennison CM, Shepherd R. Adolescent food choice: an application of the Theory of Planned Behaviour. *J Hum Nutr Diet* 1995; **8**: 9–23.
- Bryan CJ, Yeager DS, Hinojosa CP, et al. Harnessing adolescent values to motivate healthier eating. *Proc Natl Acad Sci USA* 2016; **113**: 10830–35.
- Samuels L. You and your teen and smoking. April 1, 2015. Get Healthy, Stay Healthy. <https://www.gethealthystayhealthy.com/articles/you-and-your-teen-and-smoking> (accessed Oct 26, 2021).
- Van Hook J, Quirós S, Dondero M, Altman CE. Healthy eating among Mexican immigrants: migration in childhood and time in the United States. *J Health Soc Behav* 2018; **59**: 391–410.
- Holmboe-Ottesen G, Wandel M. Changes in dietary habits after migration and consequences for health: a focus on South Asians in Europe. *Food Nutr Res* 2012; **56**: 18891.
- The Vegan Society. Why go vegan? <https://www.vegansociety.com/go-vegan/why-go-vegan> (accessed Oct 26, 2021).
- Willett W, Rockström J, Loken B, et al. Food in the Anthropocene: the EAT–*Lancet* Commission on healthy diets from sustainable food systems. *Lancet* 2019; **393**: 447–92.
- Bundy DAP, deSilva N, Horton S, Jamison D, Patton GC. Child and adolescent health and development: optimizing health outcomes. Washington, DC: World Bank, 2018.
- WHO. e-Library of Evidence for Nutrition Actions (eLENA). <https://www.who.int/elena/titles/en/> (accessed Oct 26, 2021).
- Lassi ZS, Moin A, Das JK, Salam RA, Bhutta ZA. Systematic review on evidence-based adolescent nutrition interventions. *Ann N Y Acad Sci* 2017; **1393**: 34–50.
- Ministry of Health And Family Welfare, Government of India. Adolescent health (RKSK). <https://nhm.gov.in/index1.php?lang=1&level=2&sublinkid=818&lid=221> (accessed Oct 26, 2021).
- Ruel MT, Quisumbing AR, Balagamwala M. Nutrition-sensitive agriculture: what have we learned so far? *Glob Food Secur* 2018; **17**: 128–53.
- Lavelle F, Benson T, Hollywood L, et al. Modern transference of domestic cooking skills. *Nutrients* 2019; **11**: 870.
- Slater J. Is cooking dead? The state of home economics food and nutrition education in a Canadian province. *Int J Consum Stud* 2013; **37**: 617–24.
- Holt K, Wooldridge NH, Story M, Sofka D, eds. Bright futures: nutrition, 3rd edn. American Academy of Pediatrics, 2011.
- Fernández-Gaxiola AC, De-Regil LM. Intermittent iron supplementation for reducing anaemia and its associated impairments in adolescent and adult menstruating women. *Cochrane Database Syst Rev* 2019; **1**: CD009218.
- Chong A, Cohen I, Field E, Nakasone E, Torero M. Iron deficiency and schooling attainment in Peru. *Am Econ J Appl Econ* 2016; **8**: 222–55.

- 24 Andretti B, Goldszmidt RB, Andrade EB. How changes in menu quality associate with subsequent expenditure on (un)healthy foods and beverages in school cafeterias: a three-year longitudinal study. *Prev Med* 2021; **146**: 106456.
- 25 Dickin KL, Hill TF, Dollahite JS. Practice-based evidence of effectiveness in an integrated nutrition and parenting education intervention for low-income parents. *J Acad Nutr Diet* 2014; **114**: 945–50.
- 26 Pérez-Ferrer C, Barrientos-Gutierrez T, Rivera-Dommarco JA, et al. Compliance with nutrition standards in Mexican schools and their effectiveness: a repeated cross-sectional study. *BMC Public Health* 2018; **18**: 1411.
- 27 Quinn EL, Johnson DB, Podrabsky M, Saelens BE, Bignell W, Krieger J. Effects of a behavioral economics intervention on food choice and food consumption in middle-school and high-school cafeterias. *Prev Chronic Dis* 2018; **15**: E91.
- 28 Olfert MD, Hagedorn RL, Clegg EN, Ackerman S, Brown C. Choice architecture in Appalachian high schools: evaluating and improving cafeteria environments. *Nutrients* 2019; **11**: E147.
- 29 Kenney EL, Barrett JL, Bleich SN, Ward ZJ, Cradock AL, Gortmaker SL. Impact of the Healthy, Hunger-Free Kids Act on obesity trends. *Health Aff (Millwood)* 2020; **39**: 1122–29.
- 30 Chen JL, Wilkosz ME. Efficacy of technology-based interventions for obesity prevention in adolescents: a systematic review. *Adolesc Health Med Ther* 2014; **5**: 159–70.
- 31 Wickham CA, Carbone ET. What's technology cooking up? A systematic review of the use of technology in adolescent food literacy programs. *Appetite* 2018; **125**: 333–44.
- 32 Vallabhan MK, Jimenez EY, Nash JL, et al. Motivational interviewing to treat adolescents with obesity: a meta-analysis. *Pediatrics* 2018; **142**: e20180733.
- 33 Mitchell TB, Amaro CM, Steele RG. Pediatric weight management interventions in primary care settings: a meta-analysis. *Health Psychol* 2016; **35**: 704–13.
- 34 Ruotsalainen H, Kyngäs H, Tammelin T, Kääriäinen M. Systematic review of physical activity and exercise interventions on body mass indices, subsequent physical activity and psychological symptoms in overweight and obese adolescents. *J Adv Nurs* 2015; **71**: 2461–77.
- 35 Moores CJ, Bell LK, Miller J, Damarell RA, Matwiejczyk L, Miller MD. A systematic review of community-based interventions for the treatment of adolescents with overweight and obesity. *Obes Rev* 2018; **19**: 698–715.
- 36 Thomason DL, Lukkahatai N, Kawi J, Connelly K, Inouye J. A systematic review of adolescent self-management and weight loss. *J Pediatr Health Care* 2016; **30**: 569–82.
- 37 Rivera RL, Maulding MK, Abbott AR, Craig BA, Eicher-Miller HA. SNAP-Ed (Supplemental Nutrition Assistance Program-Education) increases long-term food security among Indiana households with children in a randomized controlled study. *J Nutr* 2016; **146**: 2375–82.
- 38 Pozuelo JR, Stein A, Blakemore S-J, Kahn K, Dercon S, Pettifor A. Cash transfers in adolescence: a developmental perspective. *Lancet Child Adolesc Health* 2020; **4**: 177–78.
- 39 Carducci B, Oh C, Keats EC, Roth DE, Bhutta ZA. Effect of food environment interventions on anthropometric outcomes in school-aged children and adolescents in low- and middle-income countries: a systematic review and meta-analysis. *Curr Dev Nutr* 2020; **4**: nzaa098.
- 40 Powell LM, Leider J. The impact of Seattle's sweetened beverage tax on beverage prices and volume sold. *Econ Hum Biol* 2020; **37**: 100856.
- 41 Scarborough P, Adhikari V, Harrington RA, et al. Impact of the announcement and implementation of the UK Soft Drinks Industry Levy on sugar content, price, product size and number of available soft drinks in the UK, 2015–19: a controlled interrupted time series analysis. *PLoS Med* 2020; **17**: e1003025.
- 42 Pell D, Mytton O, Penney TL, et al. Changes in soft drinks purchased by British households associated with the UK soft drinks industry levy: controlled interrupted time series analysis. *BMJ* 2021; **372**: n254.
- 43 Taillie L, Reyes M, Colchero M, Popkin B, Corvalán C. An evaluation of Chile's Law of Food Labeling and Advertising on sugar-sweetened beverage purchases from 2015 to 2017: a before-and-after study. *PLoS Med* 2020; **17**: e1003015.
- 44 Holmes J, Meng Y, Meier PS, et al. Effects of minimum unit pricing for alcohol on different income and socioeconomic groups: a modelling study. *Lancet* 2014; **383**: 1655–64.
- 45 Godin K, Leatherdale ST, Elton-Marshall T. A systematic review of the effectiveness of school-based obesity prevention programmes for First Nations, Inuit and Métis youth in Canada. *Clin Obes* 2015; **5**: 103–15.
- 46 Lofton S, Julion WA, McNaughton DB, Bergren MD, Keim KS. A systematic review of literature on culturally adapted obesity prevention interventions for African American youth. *J Sch Nurs* 2016; **32**: 32–46.
- 47 Hawkes C, Russell S, Isaacs A, Rutter H, Viner R. What can be learned from the Amsterdam Healthy Weight programme to inform the policy response to obesity in England? Obesity Policy Research Unit (OPRU): Rapid response briefing paper. Dec 18, 2017.
- 48 Real Food Systems. Youth ambassador projects & campaigns. <https://www.realfoodsystems.org/projects> (accessed Oct 26, 2021).
- 49 Rudolf M, Perera R, Swanston D, Burberry J, Roberts K, Jebb S. Observational analysis of disparities in obesity in children in the UK: has Leeds bucked the trend? *Pediatr Obes* 2019; **14**: e12529.
- 50 Ottley PG, Dawkins-Lyn N, Harris C, et al. Childhood obesity declines project: an exploratory study of strategies identified in communities reporting declines. *Child Obes* 2018; **14**: S12–21.
- 51 Liu Z, Xu HM, Wen LM, et al. A systematic review and meta-analysis of the overall effects of school-based obesity prevention interventions and effect differences by intervention components. *Int J Behav Nutr Phys Act* 2019; **16**: 95.
- 52 Liu Z, Li Q, Maddison R, et al. A school-based comprehensive intervention for childhood obesity in China: a cluster randomized controlled trial. *Child Obes* 2019; **15**: 105–15.
- 53 Verguet S, Limasalle P, Chakrabarti A, et al. The broader economic value of school feeding programs in low- and middle-income countries: estimating the multi-sectoral returns to public health, human capital, social protection, and the local economy. *Front Public Health* 2020; **8**: 587046.
- 54 Rowe F, Stewart D, Somerset S. Nutrition education: towards a whole-school approach. *Health Educ* 2010; **110**: 197–208.
- 55 Jacob CM, Hardy-Johnson PL, Inskip HM, et al. A systematic review and meta-analysis of school-based interventions with health education to reduce body mass index in adolescents aged 10 to 19 years. *Int J Behav Nutr Phys Act* 2021; **18**: 1.
- 56 Shekar M, Kakietek J, Dayton Eberwein Julia, Walters D. An investment framework for nutrition: reaching the global targets for stunting, anemia, breastfeeding, and wasting. Directions in development. Washington, DC: World Bank, 2017.
- 57 Azzopardi PS, Hearn SJ, Francis KL, et al. Progress in adolescent health and wellbeing: tracking 12 headline indicators for 195 countries and territories, 1990–2016. *Lancet* 2019; **393**: 1101–18.
- 58 Petry N, Olofin I, Hurrell RF, et al. The proportion of anemia associated with iron deficiency in low, medium, and high human development index countries: a systematic analysis of national surveys. *Nutrients* 2016; **8**: E693.
- 59 Neufeld LM, Larson LM, Kurpad A, Mburu S, Martorell R, Brown KH. Hemoglobin concentration and anemia diagnosis in venous and capillary blood: biological basis and policy implications. *Ann N Y Acad Sci* 2019; **1450**: 172–89.
- 60 Rappaport AI, Karakochuk CD, Hess SY, et al. Variability in haemoglobin concentration by measurement tool and blood source: an analysis from seven countries. *J Clin Pathol* 2021; **74**: 657–63.
- 61 The World Bank. Adolescents out of school (% of lower secondary school age). <https://data.worldbank.org/indicator/SE.SEC.UNER.LO.ZS> (accessed Oct 26, 2021).
- 62 Plummer ML, Baltag V, Strong K, Dick B, Ross D. Global Accelerated Action for the Health of Adolescents (AA-HA!): guidance to support country implementation. Geneva: World Health Organization, 2017.
- 63 DeBar LL, Stevens VJ, Perrin N, et al. A primary care-based, multicomponent lifestyle intervention for overweight adolescent females. *Pediatrics* 2012; **129**: e611–20.
- 64 Kornet-van der Aa DA, Altenburg TM, van Randerad-van der Zee CH, Chinapaw MJM. The effectiveness and promising strategies of obesity prevention and treatment programmes among adolescents from disadvantaged backgrounds: a systematic review. *Obes Rev* 2017; **18**: 581–93.

- 65 Headey DD, Alderman HH. The relative caloric prices of healthy and unhealthy foods differ systematically across income levels and continents. *J Nutr* 2019; **149**: 2020–33.
- 66 Schoonover H, Muller M. Food without thought: how U.S. farm policy contributes to obesity. Minneapolis: Institute for Agriculture and Trade Policy (IATP), 2006.
- 67 Gillespie S, Poole N, van den Bold M, Bhavani RV, Dangour AD, Shetty P. Leveraging agriculture for nutrition in South Asia: what do we know, and what have we learned? *Food Policy* 2019; **82**: 3–12.
- 68 WHO. WHO welcomes industry action to align with global trans fat elimination targets. May 7, 2019. <https://www.who.int/news-room/detail/07-05-2019-who-welcomes-industry-action-to-align-with-global-trans-fat-elimination-targets> (accessed Oct 26, 2021).
- 69 Thow AM, Jones A, Schneider CH, Labonté R. Global governance of front-of-pack nutrition labelling: a qualitative analysis. *Nutrients* 2019; **11**: E268.
- 70 Niebyski ML, Redburn KA, Duhany T, Campbell NR. Healthy food subsidies and unhealthy food taxation: a systematic review of the evidence. *Nutrition* 2015; **31**: 787–95.
- 71 Popkin BM, Corvalan C, Grummer-Strawn LM. Dynamics of the double burden of malnutrition and the changing nutrition reality. *Lancet* 2020; **395**: 65–74.
- 72 Muth ND, Dietz WH, Magge SN, Johnson RK. Public policies to reduce sugary drink consumption in children and adolescents. *Pediatrics* 2019; **143**: e20190282.
- 73 Russell SJ, Croker H, Viner RM. The effect of screen advertising on children's dietary intake: a systematic review and meta-analysis. *Obes Rev* 2019; **20**: 554–68.
- 74 American Psychology Association. The impact of food advertising on childhood obesity. 2020. <https://www.apa.org/topics/kids-media/food> (accessed Oct 26, 2021).
- 75 Lapierre MA, Fleming-Milici F, Rozendaal E, McAlister AR, Castonguay J. The effect of advertising on children and adolescents. *Pediatrics* 2017; **140** (suppl 2): S152–56.
- 76 Braverman MT, Aarø LE. Adolescent smoking and exposure to tobacco marketing under a tobacco advertising ban: findings from 2 Norwegian national samples. *Am J Public Health* 2004; **94**: 1230–38.
- 77 Lohman BJ, Neppel TK, Lee Y, Diggs ON, Russell D. The association between household food insecurity and body mass index: a prospective growth curve analysis. *J Pediatr* 2018; **202**: 115–20.e1.
- 78 Dush JL. Adolescent food insecurity: a review of contextual and behavioral factors. *Public Health Nurs* 2020; **37**: 327–38.
- 79 Cordeiro LS, Wilde PE, Semu H, Levinson FJ. Household food security is inversely associated with undernutrition among adolescents from Kilosa, Tanzania. *J Nutr* 2012; **142**: 1741–47.
- 80 Masipa TS. The impact of climate change on food security in South Africa: current realities and challenges ahead. *Jamba* 2017; **9**: 411.
- 81 World Bank. The state of social safety nets. Washington, DC: World Bank, 2018.
- 82 DSD, SASSA, UNICEF. The South African Child Support Grant impact assessment: evidence from a survey of children, adolescents, and their households. Pretoria: UNICEF South Africa, 2012.
- 83 Blakemore S-J. The social brain in adolescence. *Nat Rev Neurosci* 2008; **9**: 267–77.
- 84 Sadler RC, Clark AF, Wilk P, O'Connor C, Gilliland JA. Using GPS and activity tracking to reveal the influence of adolescents' food environment exposure on junk food purchasing. *Can J Public Health* 2016; **107** (suppl 1): 5346.
- 85 Keats EC, Das JK, Salam RA, et al. Effective interventions to address maternal and child malnutrition: an update of the evidence. *Lancet Child Adolesc Health* 2021; **5**: 367–84.
- 86 Feng L, Wei DM, Lin ST, et al. Systematic review and meta-analysis of school-based obesity interventions in mainland China. *PLoS One* 2017; **12**: e0184704.
- 87 Liu Z, Gao P, Gao A-Y, et al. Effectiveness of a multifaceted intervention for prevention of obesity in primary school children in China: a cluster randomized clinical trial. *JAMA Pediatr* 2021; published online Nov 8. <https://doi.org/10.1001/jamapediatrics.2021.4375>.
- 88 Zhou S, Cheng Y, Cheng L, et al. Association between convenience stores near schools and obesity among school-aged children in Beijing, China. *BMC Public Health* 2020; **20**: 150.
- 89 Sheldon T. Whole city working against childhood obesity. *BMJ* 2018; **361**: k2534.
- 90 Ministry of Health and Family Welfare (MoHFW), Government of India, UNICEF, Population Council. Comprehensive National Nutrition Survey 2016–2018. New Delhi, 2019.
- 91 Sethi V, Lahiri A, Bhanot A, et al. Adolescents, diets and nutrition: growing well in a changing world. The Comprehensive National Nutrition Survey, thematic reports, issue 1, 2019.
- 92 Eat Right India. Eat Right School. <https://eatrightindia.gov.in/eatrightschool/> (accessed Oct 26, 2021).
- 93 Michaud-Létourneau I, Pelletier DL. Perspectives on the coordination of multisectoral nutrition in Mozambique and an emerging framework. *Food Policy* 2017; **70**: 84–97.
- 94 Webb P, Ghosh S, Shrestha R, et al. Measuring nutrition governance: an analysis of commitment, capability, and collaboration in Nepal. *Food Nutr Bull* 2016; **37** (suppl): S170–82.
- 95 Garrett JL, Kadiyala S, Kohli N. Working multisectorally to improve nutrition: global lessons and current status in India. POSHAN Policy Note 1. New Delhi: International Food Policy Research Institute, 2014.

Copyright © 2021 Elsevier Ltd. All rights reserved.